

## The $M_{BH}-\sigma_*$ relation of AGN2 and the unified model

Federica Ricci, Fabio La Franca (Univ. Roma Tre) & Francesca Onori (SRON)

S. Bianchi, A. Bongiorno, M. Brusa, F. Fiore, R. Maiolino, A. Marconi, E. Sani, F. Shankar, R. Schneider, C. Vignali

**ABSTRACT.** Type 2 active galactic nuclei (AGN) represent the majority of the AGN population. However, due to the difficulties in measuring their black hole (BH) masses, it is still unknown whether they follow the same BH mass-host galaxy scaling relations valid for quiescent galaxies and type 1 AGN. Here we present the locus of type 2 AGN having virial BH mass estimates in the  $M_{BH} - \sigma_*$  plane. Our analysis shows that the BH masses of type 2 AGN are  $\sim 0.9$  dex smaller than type 1 AGN at  $\sigma_* \sim 185 \text{ km s}^{-1}$ , regardless of the (early/late) AGN host galaxy morphology. Equivalently, type 2 AGN host galaxies have stellar velocity dispersions  $\sim 0.2$  dex higher than type 1 AGN hosts at  $M_{BH} \sim 10^7 M_\odot$ .

This result disagrees with standard AGN unification scenarios in which AGN1 and AGN2 are the same objects observed along different viewing angles with respect to a toroidal absorbing material.

In the luminosity range  $42.5 < \log L_X < 44.5$ , where the two distributions overlap, **AGN2 show on average significantly smaller FWHM than AGN1** ( $1970 \text{ km s}^{-1}$  instead of  $3400 \text{ km s}^{-1}$ ).




Fig. 2. *Center.* Black hole masses of AGN1 (blue open squares) and AGN2 (red filled circles) as a function of  $L_X$ . The black filled (open) circle shows the  $M_{BH}$  average value of the total AGN1 (AGN2) sample in the  $42.5 < \log(L_X/\text{erg s}^{-1}) < 44.5$  luminosity bin and has been plotted at the position of the average  $\log L_X$ . The right panel shows the projected distributions of  $M_{BH}$  of the AGN1 (blue dotted line) and AGN2 (red continuous line) samples. *Bottom:* Ratio between  $L_X$  and  $M_{BH}$  (plus a constant) of the AGN1 (blue open squares) and AGN2 (red filled circles) as a function of  $L_X$  (Onori et al. 2017b).

**The BH masses of AGN2 are  $\sim 0.9$  dex smaller than AGN1 at  $\sigma_* \sim 185 \text{ km s}^{-1}$** , regardless of the (early/late) AGN host galaxy morphology. Equivalently, AGN2 host galaxies have stellar velocity dispersions  $\sim 0.2$  dex higher than AGN1 hosts at  $M_{BH} \sim 10^7 M_\odot$ .

**This result disagrees with standard AGN unification scenarios** in which AGN1 and AGN2 are the same objects observed along different viewing angles with respect to a toroidal absorbing material.

### Related papers

Onori F., La Franca F., Ricci F. et al., 2017a, MNRAS, 464, 1783  
 Ricci F., La Franca F., Onori F., Bianchi S., 2017a, A&A, 598, A51  
 Onori F., Ricci F., La Franca F. et al., 2017b, MNRAS Letters, 468, L97  
 Ricci F., La Franca F. et al., 2017b, MNRAS Letters on line, arXiv:1706.06110




Fig. 1. *Top.* Distribution of  $L_X$  of the AGN2 (red continuous line) and Reverberation Mapping AGN1 control sample (blue dotted line). The red dot dashed and the blue dashed lines show the distribution of the AGN2 and AGN1 of the *Swift*/BAT 70-month catalogue. *Bottom.* Average FWHM of the BLR of the NIR lines (Pa $\beta$  and HeI) of AGN1 (blue open squares) and AGN2 (red filled circles) as a function of the intrinsic 14-195 keV luminosity,  $L_X$ . The black filled (open) circle shows the FWHM average value of the total AGN1 (AGN2) sample in the  $42.5 < \log(L_X/\text{erg s}^{-1}) < 44.5$  luminosity bin and has been plotted at the position of the average  $\log L_X$  (Onori et al. 2017b).

In order to compute the BH masses of the AGN2 we used the following relation calibrated by Ricci et al. (2017a):

$$\log\left(\frac{M_{BH}}{M_\odot}\right) = 7.75 + 2 \log\left(\frac{FWHM_{NIR}}{10^4 \text{ km s}^{-1}}\right) + 0.5 \log\left(\frac{L_{14-195 \text{ keV}}}{10^{42} \text{ erg s}^{-1}}\right) \quad (1)$$

As expected from the analysis of the FWHM distributions, it results that in the  $42.5 < \log(L_X/\text{erg s}^{-1}) < 44.5$  luminosity bin **the average  $M_{BH}$  of the AGN2 sample ( $\log(M_{BH}/M_\odot) = 7.08 \pm 0.10$ ) is  $\sim 0.5$  dex smaller than measured in the AGN1 sample ( $\log(M_{BH}/M_\odot) = 7.61 \pm 0.01$ ).**




Fig. 3. The  $M_{BH} - \sigma_*$  plane for local samples of RM AGN1 (blue open squares) and AGN2 (red filled circles), together with average  $M_{BH}$  of the RM AGN1 (dark blue) and AGN2 (dark red), computed in (not independent) bins of stellar velocity dispersion. The black open square (circle) shows the  $M_{BH}$  average value of the RM AGN1 (AGN2) sample in the  $135 < \sigma_* < 250 \text{ km s}^{-1}$  stellar velocity bin and has been plotted at the position of the average  $\sigma_*$  (Ricci et al. 2017b).